

Maximum Use Rates (Apply to all Turfgrass Uses)

- Do not apply more than 227 lb (0.5 lb active ingredient) per acre (5.6 lb per 1000 sq ft) per application.
- Do not apply more than 227 lb (0.5 lb active ingredient) per year and do not apply more than 882 lb (1.5 lb active ingredient) per acre (16.8 lb per 1000 sq ft) per year if using split or sequential applications.
- To maximize the potential for plant injury, do not make additional applications within five weeks of the previous application.
- In the state of New York, this product may be applied only by commercial applicators at no more than 227 lb (0.5 lb active ingredient) per acre (5.6 lb per 1000 sq ft) per year. Use of this product in New York and other states is prohibited.

Control of Crabgrass

- This product provides pre-emergence control of crabgrass (including the large, smooth, and southern species) when applied prior to germination and emergence of crabgrass at rates from 114 to 227 pounds of product (0.5 to 1.0 lb ai) per acre.
- This product also provides early post-emergence control of crabgrass applied in late spring or early summer under the 3-leaf stage of growth or earlier as described below. This type of pre- or post-emergence control is for use when crabgrass seedlings first become visible in the established turfgrass. The herbicidal activity of early post-emergence control is 1:1:2-4 week-long period of late (before or after) application) during which crabgrass control can be achieved.

Fall Applications

This product can also be applied in the late summer through early fall (late August through November) at use rates from 114 to 227 pounds per acre to provide control of crabgrass through the early part of the next spring. Fall application of this product may be followed by an appropriately timed application in the following spring for season-long control of crabgrass.

Wetness Considered or Suppressed

When applied pre-emergence (prior to germination) for control of crabgrass in the spring, this product will also control or suppress the following weeds:

Abutilon	Chenopodium	Common Purslane	Crabgrass	Digitaria	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed

Fall Applications

When applied pre-emergence (prior to germination) in the summer through early fall (late August through November), this product will control or suppress the following weeds in addition to providing pre-emergence control of crabgrass in the following spring.

Abutilon	Chenopodium	Common Purslane	Crabgrass	Digitaria	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed

WEED RESISTANCE MANAGEMENT

Damage to the active ingredient in this product is a Group 3 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain or develop plant resistance to Group 3 herbicides. Such resistance may develop in the weed population if these herbicides are used repeatedly in the same field. Such resistance may also develop in weeds that are not susceptible to Group 3 herbicides but may be affected by residual activity of other herbicides above the weeds. To manage weeds that are resistant to Group 3 herbicides, it may be necessary to rotate herbicides with other herbicides or to rotate from a different herbicide Group that is labeled for control of these weeds under the same cultural or mechanical practices. However, an herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides. Consult your local company representative, a local cooperative extension service, professional consultants or other qualified authorities to determine appropriate actions for treating specific resistant weeds.

SEED MANAGEMENT PRACTICES

Proactively implementing directed seed control strategies in crabgrass selection for weed persistence resistant to one or more herbicides is recommended. A directed seed management program may include the use of multiple herbicides with different modes of action and overlapping areas of application, or utilizing herbicides with different modes of action and different modes of action for seed control. Directed seed control strategies may include the use of herbicides for seed control, or the use of herbicides for seed control. Directed seed control strategies may include the use of herbicides for seed control, or the use of herbicides for seed control. Directed seed control strategies may include the use of herbicides for seed control, or the use of herbicides for seed control.

APPLY TO THE TURFGRASS WITH EQUIPMENT

This product may be applied with any type of equipment designed to apply granular herbicides. For best results, apply this product evenly and uniformly across the area to be treated. Avoid the use of equipment that tends to apply granules in narrow rows or concentrated areas. Obtain the proper application rate according to the manufacturer's directions. Label directions may require adjustment in higher or lower application rates under certain application conditions. The desired herbicide activity may be reduced or increased by adjusting the application rate. Apply this product uniformly over the treatment area. Some uniformity of application rate may be required by applying half of the total amount of product over the treatment area and then applying the remaining half in a different direction (e.g., at right angles to the previous direction). Avoid streaming, skips, or clumps during application.

FIELD APPLICATIONS

This product can also be applied in the late summer through early fall (late August through November) at use rates from 114 to 227 pounds per acre to provide control of crabgrass through the early part of the next spring. Fall application of this product may be followed by an appropriately timed application in the following spring for season-long control of crabgrass.

Wetness Considered or Suppressed

When applied pre-emergence (prior to germination) for control of crabgrass in the spring, this product will also control or suppress the following weeds:

Abutilon	Chenopodium	Common Purslane	Crabgrass	Digitaria	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed

Fall Applications

When applied pre-emergence (prior to germination) in the summer through early fall (late August through November), this product will control or suppress the following weeds in addition to providing pre-emergence control of crabgrass in the following spring.

Abutilon	Chenopodium	Common Purslane	Crabgrass	Digitaria	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed
Alfalfa	Cyperus	Cornflower	Crabgrass	Dandelion	Field Bindweed

WEED RESISTANCE MANAGEMENT

Damage to the active ingredient in this product is a Group 3 herbicide based on the mode of action classification system of the Weed Science Society of America. Any weed population may contain or develop plant resistance to Group 3 herbicides. Such resistance may develop in the weed population if these herbicides are used repeatedly in the same field. Such resistance may also develop in weeds that are not susceptible to Group 3 herbicides but may be affected by residual activity of other herbicides above the weeds. To manage weeds that are resistant to Group 3 herbicides, it may be necessary to rotate herbicides with other herbicides or to rotate from a different herbicide Group that is labeled for control of these weeds under the same cultural or mechanical practices. However, an herbicide mode of action classification by itself may not adequately address specific weeds that are resistant to specific herbicides. Consult your local company representative, a local cooperative extension service, professional consultants or other qualified authorities to determine appropriate actions for treating specific resistant weeds.